

2009 H1N1 Influenza Vaccine Dosage Chart

Manufacturer/ Trade name	Age	Dose-Presentation	Number of Doses/Interval	Route-Site
Novartis Vaccine/ FLUVIRIN®	Children (4 through 9 years)	0.5mL prefilled syringe or multidose vial	2* /approximately one month apart	Intramuscular
	Children (10 through 17 years)	0.5mL prefilled syringe or multidose vial	1	Intramuscular
	Adults (18 years and older)	0.5mL prefilled syringe or multidose vial	1	Intramuscular
Sanofi/	Children (6-35 months)	0.25 mL prefilled syringe or multidose vial	2* /approximately one month apart	Intramuscular
	Children (3 through 9 years)	0.5mL prefilled syringe or multidose vial	2* /approximately one month apart	Intramuscular
	10 years and older	0.5mL prefilled syringe or multidose vial	1	Intramuscular
	Adults	0.5mL prefilled syringe or multidose vial	1	Intramuscular
	18 years and older	0.5mL prefilled syringe or multidose vial	1	Intramuscular
CSL/ Afluria®	18 years and older	0.5mL prefilled syringe or multidose vial	1	Intramuscular
MedImmune/ FluMist®	Children (2 through 9 years)	0.2 mL Spray 0.1mL (1/2 dose) into each nostril as indicated on the syringe.	2*◇ /approximately one month apart	Intranasal
	Children, adolescents and adults (10 through 49 years)	0.2mL Spray 0.1mL (1/2 dose) into each nostril as indicated on the syringe.	1◇	Intranasal

*If different types of vaccine are used for the first and second doses, there is no need to revaccinate a child.

◇Two doses of LAIV (Seasonal and 2009H1N1) should not be co-administered and should be separated by at least one month (28 days). There is no restriction on co-administration of other injectable or oral live vaccines and either LAIV seasonal or LAIV 2009 H1N1 vaccine but IF not administered on the same day should be separated by at least one month. There are no restrictions on the co-administration of an injectable and LAIV flu vaccination (seasonal and 2009 H1N1) and no minimum interval between the dose of injectable and LAIV flu vaccinations (seasonal and 2009 H1N1).

